

MZ-TH/92-16

TTP92-17

April 1992 (first version)

ON GAUGE INVARIANCE OF BREIT-WIGNER PROPAGATORS

M. Nowakowski^{(a), 1} and A. Pilaftsis^{(b), 2}

^(a)Inst. für Theoretische Teilchenphysik, Universität Karlsruhe, 7500 Karlsruhe, *FRG*

^(b)Inst. für Physik, Johannes-Gutenberg Universität, 6500 Mainz, *FRG*

Final version to appear in *Z. Phys. C*

ABSTRACT

We present an approach to bosonic ($Z^0, W^\pm$) as well as fermionic (top-quark) Breit-Wigner propagators which is consistent with gauge invariance arguments. In particular, for the Z^0 -boson propagator we extend previous analyses and show that the part proportional to $k_\mu k_\nu / M^2$ must be modified near the resonance. We derive a mass shift which agrees with results obtained elsewhere by different methods. The modified form of a resonant heavy fermion propagator is also given.

¹ Present address: Physical Research Laboratory, Navrangpura, Ahmedabad 380 009, INDIA

² Address after 1, Oct. 1993, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, ENGLAND. E-mail address: pilaftsis@vipmza.physik.uni-mainz.de

The quest for the analytical form of the Breit-Wigner (BW) propagator consistent with gauge invariance has been the subject of numerous studies in the Standard Model (SM) [1-7]. Clearly its relevance nowadays lies mainly in extracting the Z^0 mass and width parameters from the resonant e^+e^- cross section. But the study of the $W^\pm$ and top propagator (we assume $m_t > M_W + m_b$ [8]) is not less important for instance in $e^+e^- \rightarrow W^*W^*$ [9,15]. In addition it has been shown that the complex phase in the BW propagator can induce new CP-violating phenomena [10-14] which can be sizeable in SM and hence can play an important role in search for new signals beyond SM. This applies both to bosonic [12-14] as well as fermionic [10-11] BW propagators. Note, however, that the fermionic BW propagator is of quite different nature than its bosonic counterpart [10].

Recently it has been shown that the part proportional to $k_\mu k_\nu$ tensor of the BW propagator for the $W^\pm$ boson must be modified in order to satisfy $U(1)$ gauge invariance [2]. In the present note we prove that the same modification must take place in the resonant Z^0 propagator. In fact, we get a mass shift (i.e. the difference between M_Z and M_R where M_R is the renormalized mass entering the equation $(q^2 - M_R^2 + iq^2\Gamma_Z/M_Z)^{-1}$) which is comparable with results obtained, for instance, in [5]. Our approach may not be regarded only as an extension of previous analyses [3,5,6], which do not consider the $k_\mu k_\nu$ part of the Z^0 BW propagator, but should also serve as an additional verification of them. Finally, with the help of Ward identities we derive the analytical form of the resonant top-quark propagator and point out its relevance for future experiments.

It should be stressed here that what we have in mind is not the full propagator, but only its resonant part. The "correctness" of our propagator should be understood in the context of resonant processes where the usage of the modified BW propagator leads consistently to gauge invariant results.

The commonly used expressions for bosonic and fermionic BW propagators are, respectively

$$\Delta_{\mu\nu}^V(q^2) = \frac{-g_{\mu\nu} + q_\mu q_\nu / M_V^2}{q^2 - M_V^2 + iM_V\Gamma_V} \quad (1)$$

$$S_F(q) = \frac{\not{q} + m_t}{q^2 - m_t^2 + im_t\Gamma_t} \quad (2)$$

where $V = Z^0, W^\pm$. The crucial step in deriving the aforementioned modification of (1) for $V = W^\pm$ is to prove that the following electromagnetic Ward identity [2]

$$k_\mu \bar{\Delta}^{\delta\rho}(p') \Gamma_{\rho\beta}^\mu(p, p', k) \bar{\Delta}^{\beta\alpha}(p) = e \left[\bar{\Delta}^{\delta\alpha}(p') - \bar{\Delta}^{\delta\alpha}(p) \right] \quad (3)$$

where

$$\Gamma_{\rho\beta}^\mu(p, p', k)|_{WW\gamma} = e \left[(p+k)_\rho g_\beta^\mu + (p'-k)_\beta g_\rho^\mu - (p+p')^\mu g_{\rho\beta} \right] \quad (4)$$

is the $W_{(\beta)}^+(p) \rightarrow W_{(\rho)}^+(p') + \gamma_{(\mu)}(k)$ vertex at tree level, is verified by inserting

$$\bar{\Delta}_{\mu\nu}^W(q) = \frac{-g_{\mu\nu} + q_\mu q_\nu / (M_W^2 - iM_W \Gamma_W)}{q^2 - M_W^2 + iM_W \Gamma_W} \quad (5)$$

and not the form given in (1). The authors of [2] have checked that the process [15]

$$e^+ + e^- \rightarrow W^{+*} + W^{-*} + \gamma \quad (6)$$

with the subsequent decays $W^\pm \rightarrow f\bar{f}'$ preserves $U(1)$ gauge invariance only if $\bar{\Delta}_{\mu\nu}^W(q)$ of (5) is used.

We mention that though the above considerations make use of the unitary gauge it is evident that they can be extended to an arbitrary ξ gauge. It is remarkable that in this case we have to make the effective replacement

$$M_W^2 \rightarrow M_W^2 - iM_W \Gamma_W \quad (7)$$

not only in the $q_\mu q_\nu$ terms of the propagator, but also in the vertices, for instance in the $\chi - W - \gamma$ vertex (χ is the charged Goldstone boson)

$$\Gamma^{\mu\nu}|_{\chi W\gamma} = ieM_W g_{\mu\nu} \rightarrow ie\sqrt{M_W^2 - iM_W \Gamma_W} g_{\mu\nu} \quad (8)$$

where the square root is to be taken on the first sheet. It will turn out that for the Z^0 and top propagator the Feynman rules must undergo a similar effective modification when one deals with a BW propagator in R_ξ gauges. However, since the Goldstone boson χ will always appear as intermediate non-asymptotic state, the above square root cannot survive in a real calculation near the W resonance.

Yet another way to understand the modification in (5) is to consider the process $q\bar{q}' \rightarrow Q\bar{Q}'$. In a ξ gauge there are two amplitudes at tree level (unitarized by a BW propagator) which we denote by $T_{W^+}^\xi(q\bar{q}' \rightarrow W^+ \rightarrow Q\bar{Q}')$ and $T_{\chi^+}^\xi(q\bar{q}' \rightarrow \chi^+ \rightarrow Q\bar{Q}')$. The gauge invariance can be checked, for instance, by

$$T_{W^+}^\xi + T_{\chi^+}^\xi \stackrel{!}{=} T_{W^+}^{\xi \rightarrow \infty} \quad (9)$$

This equation holds only if in $T_{W^+}^{\xi \rightarrow \infty}$ the resonant BW propagator is of the form given in (5). There are some subtleties concerning the replacement (7). This and the explicit prove of our statement we will, however, demonstrate not for the reaction $q\bar{q}' \rightarrow Q\bar{Q}'$, but for a similar process with an intermediate Z^0 instead of $W^\pm$. An immediate advantage is then that we will get a modified gauge-invariant form for the Z^0 propagator.

Consider, as before, the two amplitudes of the process $e^+e^- \rightarrow \nu_e\bar{\nu}_e$ with massive Dirac neutrinos

$$\begin{aligned} T_{Z^0}^\xi &= T_{Z^0}^\xi(e^+e^- \rightarrow Z^0 \rightarrow \nu_e\bar{\nu}_e) = T_{Z^0}^t + T_{Z^0}^{l,\xi} \\ T_{\chi^0}^\xi &= T_{\chi^0}^\xi(e^+e^- \rightarrow \chi^0 \rightarrow \nu_e\bar{\nu}_e) \end{aligned} \quad (10)$$

In (10) we have split the Z^0 amplitude in components proportional to $g^{\mu\nu}$ (denoted by t) and $k_\mu k_\nu$ terms (denoted by l). Note that it is not essential to consider massive Dirac neutrinos. This process is chosen as we can demonstrate our points easily in this case. We could equivalently consider the process, say $e^+e^- \rightarrow \mu^+\mu^-$, and for the sake of simplicity decouple the photon by putting $\sin\theta_W = 0$.

Our claim is now that in order to fulfill the equation

$$T_{Z^0}^\xi + T_{\chi^0}^\xi \stackrel{!}{=} T_{Z^0}^{\xi \rightarrow \infty} \quad (11)$$

it is necessary that the Z^0 BW propagator takes the same form as the W propagator in (5) with the obvious replacements $M_W, \Gamma_W \rightarrow M_Z, \Gamma_Z$. Furthermore in complete analogy to (7) we have to make the shift $M_Z^2 \rightarrow M_Z^2 - iM_Z\Gamma_Z$ in the vertices. To show this we introduce in the first step four different mass parameters. The first one just defines the position of the pole

$$\tilde{M}_Z^2 = M_Z^2 - iM_Z\Gamma_Z \quad (12)$$

Very often one uses also the definition

$$\tilde{M}_Z^2 = (M_Z - i\Gamma_Z/2)^2 \quad (13)$$

Both definitions are matters of convention. The only physical model independent quantity in a BW propagator is the complex pole position [3,5,6], i.e. $q_{pole}^2 = \tilde{M}_Z^2$, since this is a fundamental property of the S-matrix theory. For more details we refer the reader to [7].

The second mass parameter enters the longitudinal part of the Z^0 -propagator

$$\overline{\Delta}_{\mu\nu}^Z(q) = \frac{-g_{\mu\nu} + q_\mu q_\nu / (M'_Z)^2}{q^2 - \tilde{M}_Z^2} \quad (14)$$

The remaining masses appear in the χ^0 propagator and in the product of coupling constants of the pseudo-Goldstone boson χ^0 to fermions.

$$\begin{aligned} \Delta_{\chi^0}(q) &= \frac{1}{q^2 - \xi(M_Z'')^2} \\ g_{\chi^0\nu\bar{\nu}} \cdot g_{\chi^0 e\bar{e}} &= -\frac{e^2}{4\sin^2\theta_W \cos^2\theta_W (M_Z''')^2} \end{aligned} \quad (15)$$

It is sufficient to concentrate on the ξ -dependent part which can be cast into the form

$$\begin{aligned} T_{Z^0}^{l,\xi} + T_{\chi^0}^\xi &\propto \frac{1 - \xi}{q^2 - \tilde{M}_Z^2} \frac{1}{q^2 - \xi(M_Z')^2} + \frac{1}{(M_Z''')^2} \frac{1}{q^2 - \xi(M_Z'')^2} \\ &= \frac{1}{(M_Z''')^2} \left(\frac{1}{q^2 - \xi(M_Z'')^2} - \frac{1}{q^2 - \xi(M_Z')^2} \right) \\ &+ \frac{1}{(M_Z''')^2} \frac{1}{q^2 - \tilde{M}_Z^2} \frac{q^2 - \xi(M_Z''')^2 - (\tilde{M}_Z^2 - (M_Z'')^2)}{q^2 - \xi(M_Z')^2} \end{aligned} \quad (16)$$

The first term in (16) is independent of ξ only if $M_Z'' = M_Z'$. It then follows that

$$\tilde{M}_Z^2 = (M_Z')^2 = (M_Z'')^2 = (M_Z''')^2 \quad (17)$$

which completes our proof. Equation (14) together with (17) defines the consistent gauge invariant expression for the Z^0 propagator. The $k_\mu k_\nu$ part of this propagator receives the same modification as in the case of the $W^\pm$ propagator (Eq. (5)).

We would like to draw the reader's attention to one subtle point in the derivation of the modified Z^0 propagator (Eq. (14) and (17)). It concerns the shift in the Feynman rules $M_Z^2 \rightarrow M_Z^2 - iM_Z\Gamma_Z$. Instead of (15) we could equally well write the product of the χ^0 -coupling constants as

$$g_{\chi^0\nu\bar{\nu}} \cdot g_{\chi^0 e\bar{e}} = -\frac{e^2}{4\sin^2\theta_W M_W^2} \quad (18)$$

where M_Z does not appear explicitly. It would be then impossible to show the ξ independence of the amplitude. Hence if the amplitude has a resonant behaviour we have to use the product of the Goldstone bosons couplings as given in (15). Clearly the relation $\cos\theta_W = M_W/M_Z$ remains valid for physical boson masses at the zeroth order of perturbation theory. We think that this point needs a further investigation especially for a situation where both resonant particles, $W^\pm$ and Z^0 , are involved. In this note we will not pursue this topic further. $SU(2)$ Ward identities may shed some light on this problem (see treatment of the top propagator below).

Some comments concerning the status of a BW propagator are, however, in order. Feynman rules are derived from a hermitian lagrangian and are formulated for asymptotic states. The necessity to introduce a BW propagator in order to regularize an s-channel singularity reflects to the fact that our states are not asymptotic and will always appear as intermediate ones (see also another example of a t-channel singularity at the end of the paper; this singularity appears there for the same reasons). In fact, after one performs the Dyson summation of self-energy insertions, the BW propagator corresponds to *the first term* of a Laurent series expansion [3,6] of the full propagator in terms of the gauge-invariant pole q_{pole}^2 [3,5,6]. On the same footing this leading term of this expansion induces similar effective modifications in the Goldstone-boson vertices due to Ward identities. Obviously, these vertices cannot be derived, like Feynman rules, *directly* from the Lagrangian.

In a context closer to perturbation theory [3,6] we will now confirm the modification for the $k_\mu k_\nu$ part of the propagator near the resonance as previously obtained by Eqs. (14) and (17). To make this explicit, let us write the self-energy loop of the Z^0 conveniently as

$$\Pi_{\mu\nu}(q) = g_{\mu\nu}\Pi_T(q^2) + q_\mu q_\nu \Pi_L(q^2) \quad (19)$$

For simplicity we neglect $Z^0\gamma$ mixing. The inclusion of these mixing effects [17] will, however, not invalidate the discussion given below. Then, the result of the infinite summation of self-energy graphs in an arbitrary ξ gauge leads to the following form for the (unrenormalized) Z^0 propagator:

$$\begin{aligned}\Delta_{\mu\nu} &= \left[-g_{\mu\nu}(q^2 - M_Z^2 + \Pi_T) - \frac{q_\mu q_\nu}{q^2} \left(\frac{1}{\xi}(q^2 - \xi M_Z^2) + \Pi_T - q^2 \Pi_L \right) \right]^{-1} \\ &= \frac{1}{q^2 - M_Z^2 + \Pi_T} \left(-g_{\mu\nu} + \frac{(1 - \xi - \xi \Pi_L) q_\mu q_\nu}{q^2 - \xi(M_Z^2 - \Pi_T + q^2 \Pi_L)} \right)\end{aligned}\quad (20)$$

The form (20) has a physical complex pole $\tilde{M}_Z^2$ and can be determined by solving the equation

$$\tilde{M}_Z^2 - M_Z^2 + \Pi_T(\tilde{M}_Z^2) = 0 \quad (21)$$

which is independent of the gauge parameter ξ . Setting $\tilde{M}_Z^2 = M_Z^2 - iM_Z\Gamma_Z$, we can identify M_Z and Γ_Z as the physical mass and the decay width of the Z^0 boson, respectively. By expanding (20) in Laurent series [6,7], it is easy to see that the first term of this sum takes the form of Eq. (5) with M_W and Γ_W replaced by M_Z and Γ_Z , respectively. At the end of this note, we will make some additional remarks.

To compare our Z^0 propagator with other results (for instance [5]) let us rewrite (14) in such a way that the width appears to be q^2 -dependent [18]. We will retain only terms up to $\mathcal{O}(\Gamma_Z^2)$. First we define

$$\begin{aligned}\gamma_Z &= \frac{\Gamma_Z}{M_Z} \\ (M_Z^R)^2 &= M_Z^2 - \frac{\Gamma_Z^2}{4} + \Gamma_Z^2\end{aligned}\quad (22)$$

After some straightforward algebra one obtains

$$\bar{\Delta}_{\mu\nu}^Z(q) = \frac{\left(-g_{\mu\nu} + \frac{q_\mu q_\nu}{(M_Z^R)^2} (1 + i\gamma_Z) \right) (1 + i\gamma_Z)}{q^2 - (M_Z^R)^2 + iq^2 \Gamma_Z / M_Z} \quad (23)$$

where we have used the pole position defined by (13). Recall that in (23) we have neglected terms of order $\mathcal{O}(\Gamma_Z^3)$ wherever possible i.e. we have approximated $\Gamma_Z/M' \simeq$

Γ_Z/M_Z , $M'^2 + M_Z\Gamma_Z^2/M' \simeq M'^2 + \Gamma_Z^2$, $\Gamma_Z(M_Z - M') \simeq 0$ with $M'^2 = M_Z^2 - \Gamma_Z^2/4$. Expanding again in Γ_Z/M_Z we get

$$M_Z^R = M_Z \left(1 + \frac{3}{8} \Gamma_Z^2/M_Z^2 + \dots \right) \quad (24)$$

which is exactly the result obtained in [5], just by using the BW propagator of the form (5) (similar conclusions have been reached in [2-4,6]).

If we had used (12) instead of (13) then the formula (23) would remain unchanged. The only difference would be the definition of M_Z^R , namely $(M_Z^R)^2 = M_Z^2 + \Gamma_Z^2$. This results in slightly higher value for M_Z^R ($M_Z^R = M_Z(1 + \Gamma_Z^2/2M_Z^2 + \dots)$).

Finally let us discuss the case of a resonant top propagator. Similar to the case of $W^\pm$ propagator we have at any loop order

$$k_\mu \bar{S}_t^F(q) \Gamma_\gamma^\mu(p, q) \bar{S}_t^F(p) = \frac{2}{3} e \left[\bar{S}_t^F(p) - \bar{S}_t^F(q) \right] \quad (25)$$

From (25) we can deduce that the gauge invariant form for the BW propagator which respects the Ward identity at tree level is

$$i\bar{S}_t^F(p) = \frac{i}{\not{p} - m_t + i\Gamma_t/2} = i \frac{\not{p} + m_t - i\Gamma_t/2}{p^2 - m_t^2 + im_t\Gamma_t + \Gamma_t^2/4} \quad (26)$$

and not the expression given in (2). To appreciate the difference we mention that in some extensions of SM one can have $\Gamma_t/m_t \leq 0.1$ [11]. The discrepancy between $m_t^R = (m_t^2 - \Gamma_t^2/4)^{1/2}$ entering Eq. (2) and m_t could be then of the order of GeV . Clearly since the top physics is not expected to be as clean as the data on the Z^0 pole the use of this shift may not be so stringent. Yet another possible application of (26) are CP-violating processes. This possibility has been discussed in full detail in [10-11].

It is worth noting that inserting the wrong propagator (2) in the amplitude for the process $W^+d \rightarrow W^+bg$, for example, would violate the gauge symmetry even while making $SU(3)_C$ gauge transformation of the gluonic field.

At the end of the discussion of the top propagator we again address the question on the modification of Feynman rules analogous to (7). The gauge transformations

of the $W^\pm$ and Z^0 fields read

$$\begin{aligned} W_\mu^+ &\rightarrow W_\mu^+ + i \frac{\partial_\mu \chi^+}{M_W} \\ Z_\mu^0 &\rightarrow Z_\mu^0 + \frac{\partial_\mu \chi^0}{M_Z} \end{aligned} \quad (27)$$

In terms of Ward identities we then get

$$\begin{aligned} &\frac{k_\mu}{M_W} i\bar{S}_t^F(\not{p}_t) i\Gamma_{W^+}^\mu(p_t, p_D) iS_D^F(\not{p}_D) \\ &+ i\bar{S}_t^F(\not{p}_t) i\tilde{\Gamma}_{\chi^+}(p_t, p_D) iS_D^F(\not{p}_D) \\ &= \frac{g_W}{2\sqrt{2}M_W} V_{tD} \left[i\bar{S}_t^F(\not{p}_t)(1 + \gamma_5) - (1 - \gamma_5)iS_D^F(\not{p}_D) \right] \end{aligned} \quad (28)$$

for the $W^+(k) + D(p_D) \rightarrow t(p_t)$ vertex and

$$\begin{aligned} &-i \frac{k_\mu}{M_Z} i\bar{S}_t^F(\not{p}) i\Gamma_{Z^0}^\mu(p, q) i\bar{S}_t^F(\not{q}) \\ &+ i\bar{S}_t^F(\not{p}) \tilde{\Gamma}_{\chi^0}(p, q) i\bar{S}_t^F(\not{q}) \\ &= -\frac{ig_W}{2M_W} \left[(g_V^t + T_3^t \gamma_5) i\bar{S}_t^F(\not{q}) + i\bar{S}_t^F(\not{p}) (g_V^t - T_3^t \gamma_5) \right] \end{aligned} \quad (29)$$

for the $Z^0(k) + t(q) \rightarrow t(p)$ vertex. In the above D denotes down-type quark, T_3^t is the weak isospin of the top quark ($T_3^t = 1/2$) and g_V^t is the vector part of the $Z^0 - t - t$ coupling (i.e. $g_V^t = 1/2 - 4\sin^2\theta_W/3$). The unphysical Goldstone boson vertices at tree level are in our case

$$\begin{aligned} \tilde{\Gamma}_{\chi^+} &= \frac{g_W}{2\sqrt{2}} V_{tD} \left[\frac{m_t - i\Gamma_t/2}{M_W} (1 - \gamma_5) - (1 + \gamma_5) \frac{m_D}{M_W} \right] \\ \tilde{\Gamma}_{\chi^0} &= -g_W T_3^t \frac{m_t - i\Gamma_t/2}{M_W} \gamma_5 \end{aligned} \quad (30)$$

This demonstrates once again that the Feynman rule must be modified consistently. In the case of a resonant top quark we have the shift $m_t \rightarrow m_t - i\Gamma_t/2$.

In summary, we have set the proper framework of the approach discussed above which defines the relevant BW propagators and the modified vertices by Eqs. (5), (8), (26), (30). The next step would be to show, with the help of the Ward identities, the gauge invariance of any resonant process. We must, however, stress again that any resonant process can always be expressed in terms of three gauge invariant quantities

[6,7], i.e. the complex pole position of the BW propagator q_{pole}^2 , the residue of the pole R_{pole} and a q^2 -dependent background term C_{back} which has no poles at $q^2 = q_{pole}^2$. In general, these three parameters can be perturbatively expanded in different orders of the relevant coupling constant. To make contact with other analyses we emphasize that *our considerations can give informations only for the quantities q_{pole}^2 and R_{pole}* (i.e. $C_{back} = 0$). However, our approach which is also based on gauge invariance arguments not only modifies the $k_\mu k_\nu$ part of the propagator, but can also be viewed as an additional justification of previous works [3,4,6]. In this context an example would be to check the R_ξ gauge invariance of the amplitude $e^+e^- \rightarrow \mu^+\mu^-$ at any electroweak loop level without neglecting the fermion masses, in a manner similar to that presented in [6] for massless fermions.

We conclude with a general remark on Born level singularities in the kinematically allowed region. Beside the well-known infrared and s-channel singularities (e.g., like the Z^0 case) there are also t-channel singularities in the SM. Let us briefly outline what is meant by these singularities. The t-channel singular processes are closely connected to sattering processes with unstable particles in the initial state. Consider for example the scattering $\nu_e W^+ \rightarrow \nu_e W^+$. The total amplitude at Born level is the sum of a s-channel and t-channel diagram.

$$\begin{aligned} A(\nu_e W^+ \rightarrow \nu_e W^+) &= A_s(s, t) + A_t(s, t) \\ A_t(s, t) &= \frac{1}{t - m_e^2} f(s, t) \end{aligned} \tag{31}$$

where m_e is the electron mass and $f(s, t)$ is a kinematical function. The physical boundary can be easily calculated to be

$$\begin{aligned} t_{max} &= \frac{M_W^4}{s} \\ t_{min} &= 2M_W^2 - s \end{aligned} \tag{32}$$

It is clear that for

$$2M_W^2 - m_e^2 \leq s \leq \frac{M_W^2}{m_e^2} M_W^2 \tag{33}$$

the value $t = m_e^2$ lies within the physical region and one picks up a singularity in $A_t(s, t)$. There are of course many other examples of t-channel singular amplitudes, even in $2 \rightarrow 3$ processes like $W^+ d \rightarrow W^+ b g$ [19]. One of the common feature of these

singular processes is their non-localization in space-time. This can be seen as follows. At a time $t = t_0$ the initial $W_{(i)}^+$ decays into $e^+ + \nu_e^{(f)}$ (with a disconnected $\nu_e^{(i)}$). At $t = t_1 > t_0$ e^+ recombines with $\nu_e^{(i)}$ to produce $W_{(f)}^+$ (with a disconnected $\nu_e^{(f)}$ which has already been produced in the first step).

The reason for the catastrophic behaviour of $A_t(s, t)$ can be traced partly back to the fact that unstable particles cannot be prepared as asymptotic states. This is an old well-known problem [16,20]. However, we cannot get rid of the t-channel singularity when we replace the troublesome $W_{(i)}^+$ by the well-defined asymptotic state $e^+\bar{\nu}_e$ and consider instead $3 \rightarrow 2$ or $3 \rightarrow 3$ processes like $e^+\bar{\nu}_e\nu_e \rightarrow \nu_e W^+$. This problem is not new and has been a subject of numerous investigations in the days of S -matrix theory [21]. In the case of the $3 \rightarrow 2$ or $3 \rightarrow 3$ processes *this t-channel singularity cannot be cured even by introducing a BW propagator*.

The phenomenological relevance of the t-channel singular processes is quite limited (unless one insists on using the Equivalent Vector Boson Approximation [22]). However, since they belong to the class of singularities which appear at Born level they are important from theoretical point of view. The point we would like to make is that the common origin of all tree level singularities is the truncation of the Feynman-Dyson expansion at certain order. The subtle point is that this perturbation expansion can be shown to be unitary as a whole i.e. in the form

$$S = 1 + i \sum_{n=1}^{\infty} T^{(n)} \quad (34)$$

but for a finite number of terms in (34) this is, *in general*, not the case [23].

Note added. After our manuscript has been submitted we became aware of two papers [24], [25] which treat similar subject. Ref. [24] arrives in part at similar conclusion. The authors of [25] seem, however, to have obtained a different modification of the $q_\mu q_\nu$ terms in the propagator. We would like to make some comments on their

result. Ref. [25] uses the following decomposition of the self-energy tensor $\Pi_{\mu\nu}$

$$\Pi_{\mu\nu}(q) = \left(-g_{\mu\nu} + \frac{q_\mu q_\nu}{q^2} \right) R_T + \frac{q_\mu q_\nu}{q^2} R_L \quad (35)$$

After summing the absorptive parts of R_T and R_L ($\text{Im}R_T = \epsilon_T$, $\text{Im}R_L = \epsilon_L$), they obtain the following form for the propagator in the unitary gauge:

$$D_{\mu\nu} = \frac{1}{q^2 - M_W^2 + i\epsilon_T} \left[-g_{\mu\nu} + \frac{q_\mu q_\nu (q^2 + i\epsilon_T + i\epsilon_L)}{q^2 (M_W^2 + i\epsilon_L)} \right] \quad (36)$$

However, some difficulties in using such an expression in practical calculations of transition matrix elements to a given order of perturbation theory may indeed arise. The reason is that the functions $\epsilon_T(q^2)$ and $\epsilon_L(q^2)$ are generally ξ -dependent quantities in the off-resonance region [3,4], as well as the derivative $\epsilon'_T(M_W^2)$ [4], and hence the propagator (36) induces ξ -(gauge) dependence to all orders. On the other hand, ξ -terms coming from the vertices can remove the gauge dependence only to a given power of coupling constants, leaving matrix element gauge *non*-invariant at higher orders. Furthermore, it has been stressed by Sirlin [3] and Stuart [6] that the complex pole position of the propagator $\tilde{M}^2 = M^2 - iM\Gamma$ is the right expansion parameter since it is intrinsically gauge invariant. The Laurent expansion in $\tilde{M}^2$ is also the right procedure to obtain the BW propagator from the full one [6]. From Eq. (20) (and using (35)) one then gets in the unitary gauge

$$\Delta_{\mu\nu} = \frac{\text{Res}\Delta_{\mu\nu}}{q^2 - \tilde{M}^2} + \dots = Z \left[\frac{-g_{\mu\nu} + q_\mu q_\nu / \tilde{M}^2}{q^2 - \tilde{M}^2} + \frac{q_\mu q_\nu}{M^4} \left(R'_T - \frac{R_T + R_L}{M^2} \right) \right] + \dots \quad (37)$$

where $Z = [1 + R'_T(M^2)]^{-1}$ plays the role of the wave-function renormalization constant. The second term of the r.h.s. of Eq. (37) is a non-resonant background term of $\mathcal{O}(\alpha)$, while the ellipses denote contributions of $\mathcal{O}(\alpha(q^2 - \tilde{M}^2)/q^2)$. We consider this as an additional check that both approaches are consistent. Also, both propagators (Eq. (36) and Eq. (37)) obey the optical theorem as has been explicitly demonstrated in [25] for the resonant point $q^2 = M^2$.

Acknowledgements. We gratefully acknowledge valuable discussions with R.Decker, Z.Was, K. Schilcher, J. G. Körner and M. Lavelle. We also wish to thank

D. Atwood, G. Eilam, R. Mendel, R. Migneron and A. Soni for useful comments and criticisms. The work of A.P. has been supported by a grant from the Postdoctoral Graduate College of Mainz and the work of M.N. by Bundesministerium für Forschung und Technologie under the Grant Nr. 06KA757.

REFERENCES

- [1] R.G. Stuart, in Z^0 physics, Proc. XXVth Rencontres de Moriond, ed. J.T.T. Van (editions Frontières, Gif-sur-Yvette, 1990), p. 41; M. Consoli, A. Sirlin, in Physics at LEP, eds J. Ellis and R. Peccei (CERN-86-02, 1986) Vol. 1, p. 63ff; B. W. Lynn et al., *ibid.*, p. 115ff; D.C. Kenedy, B.W. Lynn, C.J.-C. Im, R.G. Stuart, Nucl. Phys. **B321** (1989) 83; A. Borrelli, M. Consoli, L. Maiani and R. Sisto, Nucl. Phys. **B333** (1990) 357; J. Ellis, ‘Status of Electroweak Interactions’, Joint Inter. Lepton- Photon Symposium and Europhysics Conference on High Energy Physics, Geneva, (1991).
- [2] G. Lopez Castro, J. L. M. Lucio, J. Pestieau, Mod. Phys. Lett. **A6** (1992) 3679 and references therein.
- [3] A. Sirlin, Phys. Rev. Lett. **67** (1991) 2127.
- [4] A. Sirlin, Phys. Lett. **B267** (1991) 240.
- [5] S. Willenbrock, G. Valencia, Phys. Lett. **B259** (1991) 373.
- [6] R. G. Stuart, Phys. Lett. **B262** (1991) 113; Phys. Lett. **B272** (1991) 353
- [7] R. G. Stuart, CERN-preprint, CERN-TH. 6261/91, Workshop on High-Energy Phenomenology, Mexico City, Mexico 1991 (QCD161, W568, 1991), and references therein.
- [8] The CDF Collaboration, F. Abe et al., Phys. Rev. **D45** (1992) 3921.
- [9] J. Layssac, G. Moulhaca, F. M. Renard and G. Gounaris, preprint of Montpellier University in 1991, PM/90-42.
- [10] A. Pilaftsis, Z. Phys. **C47** (1990) 95.

- [11] M. Nowakowski, A. Pilaftsis, Mod. Phys. Lett. **A6** (1991) 1933.
- [12] A. Pilaftsis, M. Nowakowski, Phys. Lett. **B245** (1990) 185.
- [13] G. Eilam, J. L. Hewett, A. Soni, Phys. Rev. Lett. **67** (1991) 1979; **68** (1992) 2103 (Comment); R. Cruz, B. Grzadkowski, J.F. Gunion, Phys. Lett. **B289** (1992) 440.
- [14] A. S. Joshipura, S. D. Rindani, Phys. Rev. **D46** (1992) 3008; D. Atwood et al., Phys. Rev. Lett. **70** (1993) 1364.
- [15] A. Aepli, D. Wyler, Phys. Lett. **B262** (1991) 125.
- [16] M. Veltman, Physica **29** (1963) 186.
- [17] L. Baulieu and R. Coquereaux, Ann. Phys. **140** (1982) 163.
- [18] D.Y. Bardin, A. Leike, T. Riemann, M. Sachwitz, Phys. Lett. **B206** (1988) 539.
- [19] M. Nowakowski, A. Pilaftsis, Mod. Phys. Lett **A4** (1989) 821; Z. Phys. **C42** (1990) 449.
- [20] J. Schwinger, Ann. Phys.(N.Y.) **9** (1960) 169.
- [21] R. J. Eden, P. V. Landshoff, P. J. Olive, J. C. Polkinghorne, ‘The Analytic S-Matrix’, Cambridge University Press, Cambridge 1966.
- [22] S. Dawson, Nucl. Phys. **B249** (1984) 42; R. M. Godbole and S. D. Rindani, Phys. Lett. **B190** (1987) 192.
- [23] S. N. Gupta, ‘Quantum Electrodynamics’, Gordon and Breach 1977, p.192.

[24] J. Liu, Philadelphia preprint UPR-0525T (Sept. 1992), Phys. Rev. **D47** (1993) R1741.

[25] D. Atwood et al., Technion preprint 1992, TECHNION-PH-92-39.